

✧ RESEARCH PAPER ✧

# *Breast-feeding and human immunodeficiency virus infection: Assessment of knowledge among clinicians in Kenya*

**Florence Murila MBChB MMed (Paeds)**

*Senior Lecturer, Department of Pediatrics, School of Medicine, University of Nairobi, Nairobi, Kenya*

**Moses M Obimbo MBChB MSc**

*Lecturer, Department of Human Anatomy, School of Medicine, University of Nairobi, Nairobi, Kenya*

**Rachel Musoke MBChB MMed (Paeds)**

*Professor, Department of Pediatrics, School of Medicine, University of Nairobi, Nairobi, Kenya*

**Isaac Tsikhutsu MBChB MMed (Paeds)**

*Paediatrician, Division of Child Health, Ministry of Health, Nairobi, Kenya*

**Santau Migiro MBChB MMed (Paeds)**

*Paediatrician, Division of Child Health, Ministry of Health, Nairobi, Kenya*

**Julius Ogeng'o BSc MBChB PhD**

*Professor, Department of Human Anatomy, School of Medicine, University of Nairobi, Nairobi, Kenya*

*Accepted for publication April 2013*

Murila F, Obimbo MM, Musoke R, Tsikhutsu I, Migiro S, Ogeng'o J. *International Journal of Nursing Practice* 2015; 21: 37–42

## **Breast-feeding and human immunodeficiency virus infection: Assessment of knowledge among clinicians in Kenya**

In Kenya, human immunodeficiency virus (HIV) prevalence ranks among the highest in the world. Approximately 60 000 infections yearly are attributed to vertical transmission including the process of labour and breast-feeding. The vast of the population affected is in the developing world. Clinical officers and nurses play an important role in provision of primary health care to antenatal and postnatal mothers. There are a few studies that have explored the clinicians' knowledge on breast-feeding in the face of HIV and in relation to vertical transmission this being a vital component in prevention of maternal-to-child transmission. The aim of this study was to evaluate clinicians' knowledge on HIV in relation to breast-feeding in Kenya. A cross-sectional survey was conducted to assess knowledge of 161 clinical officers and nurses serving in the maternity and children' wards in various hospitals in Kenya. The participants were derived from all district and provincial referral facilities in Kenya. A preformatted questionnaire containing a series of questions on HIV and breast-feeding was administered to clinicians who were then scored and analyzed. All the 161 participants responded.

*Correspondence: Moses M. Obimbo, School of Medicine, University of Nairobi, P.O. Box 30197, 00100 Nairobi, Kenya. Email: moses.obimbo@uonbi.ac.ke*

Majority of clinicians (92%) were knowledgeable regarding prevention of mother-to-child transmission. Regarding HIV and breast-feeding, 49.7% thought expressed breast milk from HIV-positive mothers should be heated before being given. Majority (78.3%) thought breast milk should be given regardless of availability of alternatives. According to 74.5% of the participants, exclusive breast-feeding increased chances of HIV transmission. Two-thirds (66.5%) would recommend breast-feeding for mothers who do not know their HIV status (66.5%). This study observes that a majority of the clinicians have inadequate knowledge on breast-feeding in the face of HIV. There is need to promote training programmes on breast-feeding and transmission of HIV from mother to child. This can be done as in-service training, continuous medical education and as part of the formal training within medical institutions.

**Key words:** breast-feeding, HIV, Kenya, transmission.

## INTRODUCTION

The prevalence of human immunodeficiency virus (HIV) in Kenya ranks among the highest in the world.<sup>1,2</sup> It is estimated that approximately 60 000 new infections yearly are attributable to vertical transmission that occurs during pregnancy, labour, delivery or breast-feeding.<sup>3-5</sup> Worldwide, about one-third to one-half of vertical transmission of HIV is related to breast-feeding.<sup>4,6</sup> Developing countries are grappling with an estimated 95% of total HIV infection among women and bear the greatest burden of the disease.<sup>4</sup> Armed with interventional measures, developed countries have been able to reduce the rate of vertical transmission. Consequently, focus should be shifted towards mitigating the spread of HIV through vertical transmission in developing countries.

Although trained HIV/AIDS counsellors and nutritionists offer good auxiliary efforts towards achieving high-quality prevention of maternal-to-child transmission (PMTCT) services and decrease the workload on full-time clinicians, the latter still play an unrivalled role in provision of these services.<sup>7,8</sup> There are few studies that have evaluated clinicians' knowledge on breast-feeding in the presence of HIV and in relation to vertical transmission, this being a vital component in PMTCT. This study aimed to assess clinicians' knowledge on breast-feeding in the presence of HIV.

## METHODS

A cross-sectional survey was conducted to assess the knowledge of 161 nurses and clinical officers serving in maternity and children's wards. The participants were derived from all district and provincial referral facilities in Kenya in the year 2010 and attending a seminar on essentials of newborn care. Authority to carry out the study was granted by the Ministry of Public Health and Sanitation. Professional nurses are those with 3 years of training at a medical training college, whereas clinical officers have training in clinical medicine, also at a medical training

college, and work closely with medical officers who are university graduates.

Closed-ended questions on HIV transmission and preventive measures that were of special relevance to prevention of mother-to-child transmission were used in order to assess the working knowledge of the interviewees. The questionnaires were pretested by a consultant paediatrician working in the newborn unit through participation in a pretest by the health-care providers within the unit at Kenyatta National Hospital, and adjustments were made accordingly. Questions included the aspects of initiation of breast-feeding, risks of not breast-feeding, breast-feeding after 6 months and risks of vertical transmission. Responses were compared with those recommended by WHO/UNAIDS guidelines. The data obtained were entered and analyzed with Statistical Package for the Social Sciences 16.0 (SPSS Inc., Chicago, IL, USA). Frequencies and means were obtained for various parameters, whereas the  $\chi^2$ -test was used to test the difference in knowledge between the clinical officers and the nurses.

## RESULTS

All 161 participants who attended the seminar responded. The mean age was 29.1 years, with 91 females and 70 males. One hundred and six (65.8%) were nurses and 55 (34.2%) clinical officers (Table 1). The majority of the

**Table 1** General characteristics of respondents

	Nurse s (n = 106)	Clinical officers s (n = 55)
Male (%)	36	81
Mean age (years)	33.6	24.3
With training on PMTCT (%)	96	88

PMTCT, prevention of mother-to-child transmission.

**Table 2** Respondents' answers to questions on breast-feeding practices

	Correct responses, <i>n</i> (%)			<i>P</i> value <sup>†</sup>
	Nurses ( <i>n</i> = 106)	Clinical officers ( <i>n</i> = 55)	Total	
For mothers who are HIV-positive:				
Should breast milk be heated before being given to infants?	58 (54.7)	22 (40)	80 (49.7)	0.295
In the presence of alternative feeding programmes, should breast milk be given?	20 (19)	15 (27.3)	35 (21.7)	0.572
Should breast-feeding continue after weaning at 6 months?	94 (88.6)	42 (39.6)	136 (84.5)	0.088
Does exclusive breast-feeding for 6 months increase chances of HIV transmission?	16 (15.1)	25 (45.5)	41 (25.5)	<0.001
Should breast-feeding be encouraged for mothers who don't know their HIV status?	83 (78.3)	24 (43.6)	105 (65.2)	<0.05
Does lack of breast-feeding increase risk of childhood illnesses and death?	78 (73.5)	43 (78.1)	121 (75.2)	0.708
Does mastitis increase chances of HIV transmission?	60 (56.6)	40 (72.7)	100 (62.1)	0.205
Mean	58 (54.7)	30 (54.5)	88 (54.7)	—

Questions based on the WHO/UNAIDS breast-feeding guidelines for HIV-infected mothers. <sup>†</sup> $\chi^2$ -test.

clinicians (92%) were knowledgeable regarding the PMTCT programme and what it entails.

Regarding HIV and breast-feeding, when asked whether breast-feeding is recommended for mothers who do not know their HIV status, 66.5% answered in the affirmative; on the subject of the infant feeding at the breast, 49.7% of the respondents thought expressed breast milk from HIV-positive mothers should be heated before being given, 78.3% thought breast milk from HIV-positive mothers should be given regardless of the availability of alternatives, and 75.2% of the clinicians conceded that the risk of childhood illnesses and death outweighed the risk of HIV transmission from breast milk if breast-feeding was not done. Asked whether breast-feeding should continue after weaning at 6 months, 84.5% affirmed that it should be continued. On the risk of vertical transmission, 74.5% responded that exclusive breast-feeding for 6 months increased the chances of HIV transmission as opposed to giving additional feeding. Moreover, 38% failed to relate the contribution of mastitis to increased risk of HIV transmission (Table 2).

## DISCUSSION

This study observes discrepant knowledge levels about breast-feeding practices in those who are supposed to take

care of mothers infected with HIV in Kenya. This is despite the fact that breast-feeding is known to be the best method of infant feeding and should be encouraged exclusively for the first 6 months of an infant's life.<sup>9–12</sup>

Less than 50% of respondents would recommend heating of breast milk before feeding the baby of an HIV-infected mother. Despite WHO recommendations that heating provides a simple yet effective treatment method to inactivate cell-free HIV,<sup>13</sup> the clinicians in this study seemed not to be aware of this concept. An earlier study conducted in Kenya showed that most health informants did not believe that heat treatment of milk could destroy HIV; hence, they similarly did not favor this option.<sup>14</sup> The various methods of heat treatment are effective and can easily be performed at home. They retain the nutritional potency and immunoglobulin content of breast milk in addition to being superior to feeding with formula milk.<sup>15,16</sup> Obstacles that are faced in using these methods, including social stigma, time and physical constraints, can easily be addressed through family and community support. The clinicians should be supported to learn these methods accordingly.

Although most of the clinicians had heard of PMTCT, only 74% knew about the risk of postnatal transmission of HIV via breast milk. This is in contrast to earlier

observations among community members and health workers, where a larger percentage (85.5%) of the respondents knew about HIV transmission through breast milk.<sup>14</sup> In general, nurses were the least knowledgeable cadre ( $P < 0.001$ ). Nurses are the health professionals who are in regular contact with mothers and their infants. They live in the local communities and are in a strategic position to encourage exclusive breast-feeding in hospitals and at home.<sup>17,18</sup>

The majority (84.9%) believed that exclusive breast-feeding increased chances of HIV transmission. Indeed, it has been noted that the majority of HIV-positive women in Kenya practise mixed feeding shortly after giving birth, and this has been blamed on lack of proper information.<sup>19,20</sup> Exclusive breast-feeding for 6 months followed by abrupt weaning has been proposed as a protective strategy for HIV-positive mothers in communities where formula is not affordable or safe to use.<sup>21,22</sup> There is therefore need for specific training on breast-feeding and issues related to mother-to-child transmission of HIV. The findings recorded in the current study support literature reports on the lethargic attitude and practices of health professionals towards breast-feeding, which might be due to poor training.<sup>23–25</sup> Internationally, the consensus is that when replacement or alternative feeding is acceptable, feasible, affordable, sustainable and safe (AFASS), avoidance of all breast-feeding by HIV-infected mothers should be encouraged.<sup>26</sup> Otherwise, expressed breast milk, heat treatment and cup-feeding could serve as an affordable and safe source of nutrition and immunological protection during the first 2 years of an infant's life and would help in preventing them from developing malnutrition and its attendant complications.<sup>13</sup> In Africa, a significant proportion of the population cannot access alternative feeding programmes, and many HIV-positive women choose to carry out mixed feeding.<sup>27</sup> It is imperative to equip clinicians with adequate knowledge on alternative feeding and safe breast-feeding practices for HIV-positive women living in resource-limited settings.

In Kenya, although as part of PMTCT programmes mothers are required to undergo HIV testing as part of the antenatal profile, full compliance has not been achieved.<sup>28,29</sup> It is therefore encouraging to note that in the current study many nurses affirm that they would encourage mothers to breast-feed exclusively regardless of their unknown HIV status. Pertinent to this observation are reports that exclusive breast-feeding is the best option for an infant because it protects them from morbidity and

mortality, HIV-related or not.<sup>30</sup> Indeed, HIV-exposed infants who were breast-fed exclusively for at least 3 months had a lower risk of HIV infection than mixed-fed infants.<sup>21</sup> The majority of clinical officers (56.4%), on the contrary, would not encourage the mothers to breast-feed ( $P < 0.05$ ). This suggests that further education of clinical officers is required.

A significant proportion of clinicians (43.4% of nurses and 27.3% of clinical officers) in this study could not positively relate the effect of mastitis to increased chances of HIV transmission. This result is comparable with observations made by Lawrence<sup>31</sup> that mastitis is not taken seriously by medical personnel. Mastitis is an inflammatory process in the breast and occurs in about 30% of lactating mothers.<sup>32,33</sup> In mastitis, inflammatory cells and extracellular fluid extravasate and enter the breast milk. Consequently, inflammatory cells such as infected lymphocytes could raise HIV-1 load in breast milk and raise the chances of mother-to-child transmission of HIV.<sup>34</sup> These results imply that there should be more effort geared towards increasing awareness of mastitis and its attendant complications.

To achieve informed decisions when it comes to infant care and feeding in the presence of HIV, various organizations have come up with guidelines to help clinicians. UNICEF, for example, has a programme aimed at improving infant and young child feeding at a community level. This is in recognition of the fact that many clinicians in developing countries lack the prerequisite practical support, one-to-one counseling skills and correct information. The purpose of this programme is to ensure equitable access to health care for all infants at a community level.<sup>35</sup>

A study conducted among counsellors in northern Tanzania on perspectives on HIV/AIDS and breast-feeding found that not only did patients have difficulty in understanding feeding options for HIV-infected mothers, but so did the counsellors. In their conclusion, the authors were of the view that the counsellors needed additional training in nondirective counselling and infant feeding options to better equip them.<sup>36</sup> The WHO has developed a guideline for health supervisors and managers on how to support and promote safe infant feeding practices in the context of HIV.<sup>37</sup> The guidelines are simple to follow and practical. If these guidelines are incorporated in medical school curricula and continuous medical education programmes, they are likely to promote safe infant feeding practices with better outcomes.

## CONCLUSION

This study observes that a majority of the clinicians had inadequate knowledge on breast-feeding in the presence of HIV. There is need to promote training programmes on breast-feeding and transmission of HIV from mother to child. This can be done as in-service-training, as continuous medical education and as part of formal training within medical institutions.

## REFERENCES

- 1 Joint United Nations Programme on HIV/AIDS. UNAIDS report on the global AIDS epidemic 2010. Geneva: UNAIDS, 2010.
- 2 National AIDS Control Council, Kenya. UNGASS 2010: United Nations General Assembly Special Session on HIV and AIDS. Country report—Kenya. Nairobi: National AIDS Control Council, 2010.
- 3 Colton TC. Preventing mother-to-child transmission of HIV in Kenya. Pathfinder International's experience: 2002–2005. Watertown, MA: Pathfinder International, 2005.
- 4 World Health Organization. HIV and infant feeding: a review of HIV transmission through breast-feeding. Geneva: World Health Organization, 1998. Publication Nos. WHO/FRH/NUT 98.3, UNAIDS 98.5, UNICEF/PD/NUT(J) 98.3.
- 5 Committee on Infectious Diseases, American Academy of Pediatrics. *Red Book: Report of the Committee on Infectious Diseases*, 25th edn. Elk Grove Village, IL, USA: American Academy of Pediatrics, 2000.
- 6 Cohn SE, Clark RA. Human immunodeficiency virus infection in women. In: Mandell GL, Bennett JE, Dolin R (eds). *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*, 5th edn. Philadelphia: Churchill Livingstone, 2000; 1452–1467.
- 7 Bassett MT. Ensuring a public health impact of programs to reduce HIV transmission from mothers to infants: the place of voluntary counseling and testing. *American Journal of Public Health* 2002; **92**: 347–351.
- 8 United Nations Children's Fund. Evaluation of United Nations-supported pilot projects for the prevention of mother-to-child transmission of HIV. New York: UNICEF, 2003.
- 9 World Health Organization. Global strategy for infant and young child feeding: the optimal duration of exclusive breastfeeding. Geneva: World Health Organization, 2001. 54th World Health Assembly provisional agenda item 13.1.
- 10 Kramer MS, Chalmers B, Hodnett ED *et al.* Promotion of Breastfeeding Intervention Trial (PROBIT). *JAMA: The Journal of the American Medical Association* 2001; **285**: 413–431.
- 11 Morrow AL, Guerrero ML, Shults J *et al.* Efficacy of home based peer counselling to promote exclusive breast-feeding: a randomised control trial. *Lancet* 1999; **353**: 1226–1231.
- 12 Bhandari N, Bahl R, Mazumdar S *et al.* Effect of community-based promotion of exclusive breastfeeding on diarrhoeal illness and growth: a cluster randomised controlled trial. *Lancet* 2003; **361**: 1418–1423.
- 13 Rollins N, Meda N, Becquet R *et al.* Preventing postnatal transmission of HIV-1 through breast-feeding: modifying infant feeding practices. *Journal of Acquired Immune Deficiency Syndromes* 2004; **35**: 188–195.
- 14 Wachira J, Otieno-Nyunya B, Ballidawa J, Braitstein P. Assessment of knowledge, attitudes and practices of infant feeding in the context of HIV: a case study from western Kenya. *Journal of Social Aspects of HIV/AIDS Research Alliance* 2009; **6**: 120–133.
- 15 Chantry CJ, Wiedeman J, Buehring G *et al.* Effect of flash-heat treatment on antimicrobial activity of breastmilk. *Breastfeeding Medicine* 2011; **6**: 111–116.
- 16 Israel-Ballard KA, Abrams BF, Coutsooudis A *et al.* Vitamin content of breast milk from HIV-1-infected mothers before and after flash-heat treatment. *Journal of Acquired Immune Deficiency Syndromes* 2008; **48**: 444–449.
- 17 Akinsola HA. Fostering hope in people living with AIDS in Africa: the role of primary health-care workers. *The Australian Journal of Rural Health* 2000; **9**: 158–165.
- 18 Lagner S, Hutelmyer C. Patient satisfaction with outpatient human immunodeficiency virus care as delivered by nurse practitioners and physicians. *Holistic Nursing Practice* 1995; **10**: 54–60.
- 19 Kiarie JN, Richardson BA, Mbori-Ngacha D, Nduati RW, John-Stewart GC. Infant feeding practices of women in a perinatal HIV-1 prevention study in Nairobi, Kenya. *Journal of Acquired Immune Deficiency Syndromes* 2004; **35**: 75–81.
- 20 Chopra M, Doherty T, Jackson D, Ashworth A. Preventing HIV transmission to children: quality of counselling of mothers in South Africa. *Acta Paediatrica* 2005; **94**: 357–363.
- 21 Coutsooudis A, Pillay K, Kuhn L *et al.* Method of feeding and transmission of HIV-1 from mothers to children by 15 months of age: prospective cohort study from Durban, South Africa. *AIDS* 2001; **15**: 379–387.
- 22 Ogundele MO, Coulter JB. HIV transmission through breastfeeding: problems and prevention. *Annals of Tropical Paediatrics* 2003; **23**: 91–106.
- 23 Freed GL, Clark SJ, Sorenson J *et al.* National assessment of physicians' breast-feeding knowledge, attitudes, training, and experience. *JAMA: The Journal of the American Medical Association* 1995; **273**: 472–476.
- 24 Philipp BL, Merewood A, Gerendas EJ, Bauchner H. Breastfeeding information in pediatric textbooks needs

- improvement. *Journal of Human Lactation* 2004; **28**: 206–210.
- 25 DiGirolamo AM, Grummer-Strawn LM, Fein SB. Do perceived attitudes of physicians and hospital staff affect breastfeeding decisions? *Birth* 2003; **30**: 94–100.
- 26 World Health Organization; Joint United Nations Programme on HIV/AIDS; United Nations Children's Fund; United Nations Population Fund. *HIV and Infant Feeding: Guidelines for Decision-Makers*. Geneva, Switzerland: World Health Organization, 2003.
- 27 Nduati R, John G, Mbori-Ngacha D *et al.* Effect of breastfeeding and formula feeding on transmission of HIV-1: a randomized clinical trial. *JAMA: The Journal of the American Medical Association* 2000; **283**: 1167–1174.
- 28 Turan JM, Bukusi EA, Onono M, Holzemer WL, Miller S, Cohen CR. HIV/AIDS stigma and refusal of HIV testing among pregnant women in rural Kenya: results from the MAMAS Study. *AIDS and Behavior* 2011; **15**: 1111–1120.
- 29 Katz DA, Kiarie JN, John-Stewart GC *et al.* Male perspectives on incorporating men into antenatal HIV counseling and testing. *International Journal of STD and AIDS* 2009; **20**: 765–767.
- 30 Piwoz E, Huffman S, Lusk D, Zehner E. Early breastfeeding cessation as an option for reducing postnatal transmission of HIV in Africa: issues, risks, and challenges. Washington: Academy for Educational Development, 2001.
- 31 Lawrence RA. *Breastfeeding: A Guide for the Medical Profession*. St. Louis, MO, USA: Mosby, 1994.
- 32 Thomsen AC, Espersen T, Maigaard S. Course and treatment of milk stasis, non-infectious inflammation of the breast, and infectious mastitis in nursing women. *American Journal of Obstetrics and Gynecology* 1984; **149**: 492–495.
- 33 Riordan J, Nichols F. A descriptive study of lactation mastitis in long-term breastfeeding women. *Journal of Human Lactation* 1990; **6**: 53–58.
- 34 Prentice A, Prentice AM, Lamb WH. Mastitis in rural Gambian mothers and the protection of the breast by milk antimicrobial factors. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1985; **79**: 90–95.
- 35 United Nations Children's Fund. *Community Based Infant and Young Child Feeding*. New York: UNICEF, 2012. Available from URL: [http://www.unicef.org/nutrition/index\\_58362.html](http://www.unicef.org/nutrition/index_58362.html). Accessed 3 March 2013.
- 36 Manuela M, Manongi R, Klepp K. Counsellors' perspectives on antenatal HIV testing and infant feeding dilemmas facing women with HIV in Northern Tanzania. *Reproductive Health Matters* 2002; **10**: 144–156.
- 37 World Health Organization; Joint United Nations Programme on HIV/AIDS. *HIV and Infant Feeding: A Guide for Health Care Managers and Supervisors*. Geneva: World Health Organization, 1998.