

Research Article

Hexafurcation of Basilar Artery in a Black Kenyan Population

Ogeng'o JA, Mwachaka PM*, Olabu BO, Ogeng'o NM and Gimongo J

Department of Human Anatomy, University of Nairobi-Kenya, Kenya

*Corresponding author: Mwachaka PM, Department of Human Anatomy, University of Nairobi, P.O. BOX 00100 – 30197 Nairobi, Kenya

Received: October 24, 2015; Accepted: January 18, 2016; Published: January 22, 2016

Abstract

Pattern of terminal branching of the basilar artery may influence predisposition to atherosclerosis and aneurysms. Knowledge of unusual patterns is also important in interpretation of extent of stroke and during neuroradiological and neurosurgical procedures at basilar artery termination. Two hundred brains of black Africans were studied for pattern of variant termination. We observed two cases of unusual termination: two hexafurcations. All these cases were associated with variant origin of superior cerebellar and thalamoperforator arteries. This suggests that variant termination of basilar artery may include hexafurcation consequent to unusual origin of superior cerebellar and thalamoperforator arteries. Neurosurgeons, neurophysicians, neuroradiologists should be aware of such strange terminal branching pattern if they are to avoid inadvertent vascular injury. We recommend preoperative evaluation of the region before surgery.

Keywords: Basilar artery; Hexafurcation; Africans; Atherosclerosis

Introduction

The Basilar Artery (BA), usually terminates by dividing into two posterior cerebral arteries [1,2]. Variant terminations described include double posterior cerebral arteries and common trunks for Posterior Cerebral Arteries (PCA) and Superior Cerebellar Arteries (SCA) trifurcation, quadrifurcation and penta-furcation [3-5]. Such variant branching patterns of arteries predisposes them to atherosclerosis and aneurysms [6,7], and may complicate surgery at basilar bifurcation angle and clivus [8]. The abnormalities may also alter the relationships with and compress oculomotor nerve [9]. There are, however, few reports on variant termination of the basilar artery. We recently reported up to 5 branches. In all cases of trifurcation, quadrifurcation and penta-furcation, the variation was due to duplication and rostral shift of superior cerebellar artery [4]. Potentially, more than five branches is possible but hitherto unreported in spite its importance. This study, therefore investigated the pattern of termination of the basilar artery in a black Kenyan population, for cases where there were more than five terminal branches.

Materials and Methods

Two hundred brains were examined. The arachnoids matter was gently peeled off to expose the basilar artery in its entire length. Pia matter, veins and small perforating arteries were sacrificed to clarify the pattern of termination. The termination was examined further for unusual terminal branching pattern. The arteries given off just before terminal bifurcation, which coursed laterally to ramify over the superior surface of cerebellar hemispheres, were identified as Superior Cerebellar Artery (SCA). The terminal branches which were joined to internal carotid by posterior communicating were identified as Posterior Cerebral Artery (PCA). Images of these patterns were taken with a high resolution digital camera.

Results

Unusual terminal branching of basilar artery, hexafurcation, whereby the basilar artery gave rise to six unequivocal branches was noted in two cases. In one, there were two superior cerebellar arteries arising from the same point as PCA bilaterally (Figure 1A). In the second one, the unusual rostral superior cerebellar artery was duplicated unilaterally, and one enlarged thalamoperforator arose from basilar bifurcation (Figure 1B).

Discussion

Hexafurcation of basilar artery is hitherto unreported. Such branching patterns are potential sites for development of atheromatous lesions because of complex flow patterns such as recirculation and secondary flow causing abnormal haemodynamic




Figure 1A-B: Variant termination of basilar artery in a black Kenyan population.

BA: Basilar Artery; PCA: Posterior Cerebral Artery; TPA: Thalamoperforator Arteries

A) Note bilateral duplication of superior cerebellar artery (asterisk) and narrow bifurcation. B) Note unilateral duplication of superior cerebellar artery (asterisk) and thalamoperforator artery (arrow) arising from basilar artery bifurcation. A second thalamoperforator artery (arrow) arises from PCA.

events referred to as “disturbed flow” [7,10]. In our recent account of variant termination of basilar artery, the most common cause of variations was additional SCA [4]. We discussed the importance of this pattern. In the current report, duplication SCA is still an important factor. The game changer, however, are the thalamo perforator arteries. These usually arise from the initial segment of posterior cerebral and posterior communicating arteries [11,12]. In the current study, they arose from the basilar bifurcation. This is concordant with isolated reports that 1 - 5 rostral perforators may arise from basilar termination [12,13].

These variations may be attributed to the pattern of embryological development of basilar artery. This artery initially appears as a plexus of paired anterior longitudinal neural arteries which anatomize into a single basilar artery. These variation may result from defective fusion and remodeling of these arteries [14,15].

Literature is silent on the significance of these variations. Nonetheless, variant branching like trifurcations of other arteries in the brain, say of middle cerebral arteries are favorable lodging sites for cerebral emboli with consequent ischemia of the affected region [16]. Accordingly, the unusual pattern observed in the present study implies that in case of basilar artery thrombosis, embolic occlusion of the branches is most likely to occur. Knowledge of these variations is important to enable neuroradiologists and neurosurgeons, surgeons safely diagnose as well as plan and execute interventions for disorders at basilar bifurcation. For example, execute vascular by – pass and shunting for treatment of stenosis, aneurysms and A - V malformations [17].

Conclusion

Variant termination of the basilar artery may include hexafurcation consequent to unusual origin of superior cerebellar and thalamoperforator arteries. Physicians, surgeons and radiologists should be aware of the unusual terminal branching patterns when managing posterior circulatory stroke or other posterior cranial fossa pathologies. We recommend preoperative evaluation of the region before surgery.

Acknowledgement

We are grateful to Acleus Murunga and Charles Nzioka for technical assistance and to Antonina Odock – Opiko for typing the manuscript.

References

1. Yasargil MG. Clinical Considerations, Surgery of the Intracranial Aneurysms and Results. Newyork: Thieme. 1984; 403.
2. Rhoton AL Jr. The cerebellar arteries. Neurosurgery. 2000; 47: S29-68.
3. Padmavathi G, Rajeshwari T, Niranjana Murthy KV. Study of the Variations in the Origin & Termination of Basilar Artery. Anat Karnataka - Int J. 2011; 5: 54-59.
4. Olabu B. Variant termination of basilar artery in a black Kenyan population. 2012.
5. Bala M, Kaushal S, Passi DK. Trifurcation of basilar artery. Int J Anat Var IJAV. 2013; 6: 199-200.
6. Ingebrigtsen T, Morgan MK, Faulder K, Ingebrigtsen L, Sparr T, Schirmer H. Bifurcation geometry and the presence of cerebral artery aneurysms. J Neurosurg. 2004; 101: 108-113.
7. Gessaghi VC, Raschi MA, Larretguy AE, Perazzo y CA. Influence of arterial geometry on a model for growth rate of atheromas. J Phys Conf Ser. 2007; 90: 012046.
8. Dagainar A, Kaya AH, Aydin ME, Kopuz C, Senel A, Demir MT, et al. The Superior Cerebellar Artery: Anatomic Study with Review. Neurosurg Q. 2007; 17: 235-240.
9. Uchino A, Sawada A, Takase Y, Kudo S. Variations of the superior cerebellar artery: MR angiographic demonstration. Radiat Med. 2003; 21: 235-238.
10. Buchanan JR, Kleinstreuer C, Hyun S, Truskey GA. Hemodynamics simulation and identification of susceptible sites of atherosclerotic lesion formation in a model abdominal aorta. J Biomech. 2003; 36: 1185-1196.
11. Saeki N, Rhoton AL Jr. Microsurgical anatomy of the upper basilar artery and the posterior circle of Willis. J Neurosurg. 1977; 46: 563-578.
12. Pai BS, Varma RG, Kulkarni RN, Nirmala S, Manjunath LC, Rakshith S. Microsurgical anatomy of the posterior circulation. Neurol India. 2007; 55: 31-41.
13. Sakata S. Microsurgical Anatomy of the Basilar Artery: Surgical Approaches to the Basilar Trunk and Vertebrobasilar Junction Aneurysms. Korean J Cerebrovasc Dis. 2001; 3: 5-10.
14. PADGET DH. The Development of the Cranial Arteries in the Human Embryo... With Five Plates, Etc. 1948.
15. Brassier G, Morandi X, Fournier D, Velut S, Mercier P. Origin of the perforating arteries of the interpeduncular fossa in relation to the termination of the basilar artery. Interv Neuroradiol J Peritherapeutic Neuroradiol Surg Proc Relat Neurosci. 1998; 4: 109-120.
16. Futrell N. Pathophysiology of acute ischemic stroke: new concepts in cerebral embolism. Cerebrovasc Dis. 1998; 8: 2-5.
17. Mamatha H, D'Souza AS, Pallavi, Suhani S. Human cadaveric study of the morphology of the basilar artery. Singapore Med J. 2012; 53: 760-763.